



**UNIVERSITI KUALA LUMPUR  
Malaysia France Institute**

---

**FINAL EXAMINATION  
JANUARY 2010 SESSION**

---

**SUBJECT CODE** : FKB 15103  
**SUBJECT TITLE** : ENGINEERING MATHEMATICS 1  
**LEVEL** : BACHELOR  
**TIME / DURATION** : 4.00pm – 6.00pm  
( 2 HOURS )  
**DATE** : 27 APRIL 2010

---

**INSTRUCTIONS TO CANDIDATES**

---

1. Please read the instructions given in the question paper **CAREFULLY**.
2. This question paper is printed on both sides of the paper.
3. Please write your answers on the answer booklet provided.
4. Answer should be written in blue or black ink except for sketching, graphic and illustration.
5. This questions paper consists of **SIX (6)** questions. Answer **FOUR (4)** questions only.
6. Answer **ALL** questions in English.

---

**THERE ARE 4 PAGES OF QUESTIONS AND 7 PAGES OF APPENDIX, EXCLUDING THIS PAGE.**

---

(Total: 80 marks)

INSTRUCTION: Answer only FOUR questions.

Please use the answer booklet provided.

Question 1

The two points  $(-2,4)$  and  $(1,3)$  lie on a circle with equation in form  $x^2 + y^2 + ax + by + c = 0$  where  $a, b$  and  $c$  are constants.

- (a) Find two equations in  $a, b$  and  $c$  and solve the system of equations by using elementary row operations.

(10 marks)

- (b) If  $(2,2)$  is also lie on the circle, determine the value of  $a, b$  and  $c$  by using Cramer's rule. Hence, find the equation of the circle.

(10 marks)

Question 2

Given that  $(4 + j)$  is a root of the equation  $x^3 - 6x^2 + x + 34 = 0$ ,

- (a) Factorize the cubic expression  $x^3 - 6x^2 + x + 34$  completely in complex domain.

(8 marks)

- (b) Hence, decompose in complex domain the following fraction,

$$\frac{1}{x^3 - 6x^2 + x + 34}$$

(12 marks)

**Question 3**

Given transfer function of a simple amplifier when negative feedback,  $\beta$ , is applied is given by

$$T = \frac{A_0 / (1 + j\omega T_1)}{1 + \frac{A_0}{1 + j\omega T_1} \beta (1 + j\omega T_2)}$$

where  $A_0$  is the low frequency gain,  $\omega$  is the angular frequency,  $T_1$  is the amplifier time constant,  $T =$  new transfer function and  $T_2 =$  feedback time constant.

(a) Show that 
$$T = \frac{A_0}{1 + A_0\beta + j\omega(T_1 + A_0\beta T_2)}$$

(3 marks)

(b) Find the gain (modulus of  $T$ ) and phase (argument of  $T$ ) of the  $T$  at an angular frequency  $\omega = 2 \times 10^3$  rad/s,  $A_0 = 1000$ ,  $\beta = 0.1$ ,  $T_1 = 0.5 \times 10^{-3}$  s and  $T_2 = 1 \times 10^{-4}$  s.

(7 marks)

(c) A cable has the following constants :  $R = 10 \Omega$ ,  $L = 0.1 \times 10^{-3}$  H,  $G = 1 \times 10^{-6}$  siemen and  $C = 1 \times 10^{-9}$  F. For  $\omega = 10,000$  rad/s, determine the characteristic impedance,

$$Z_0 \text{ where } Z_0 = \sqrt{\frac{R + j\omega L}{G + j\omega C}}$$

(10 marks)

**Question 4**

Given a point A with coordinates (3, -1, 5) and a line  $l$  with the equation

$$\underset{\sim}{r} = \underset{\sim}{8} \underset{\sim}{i} - \underset{\sim}{k} + \lambda \left( \underset{\sim}{-6} \underset{\sim}{i} + \underset{\sim}{j} + \underset{\sim}{4} \underset{\sim}{k} \right).$$

- (a) Find the coordinates of a point B on line  $l$  if AB is perpendicular to line  $l$ .

(7 marks)

Given a plane  $\pi_1$  with the equation  $\underset{\sim}{r} \cdot \left( \underset{\sim}{i} - \underset{\sim}{j} + \underset{\sim}{3} \underset{\sim}{k} \right) = 15$ .

- (b) Find the coordinates of a point C on  $l$  if the line  $l$  intersects the plane  $\pi_1$ .

(5 marks)

- (c) Determine the equation of a plane  $\pi_2$  which contains the points A, B and C.

(8 marks)

**Question 5**

- (a) If  $y = \ln(\sin px + \cos px)$ , where  $p$  is a constant, show that

$$\frac{d^2 y}{dx^2} + \left( \frac{dy}{dx} \right)^2 + p^2 = 0.$$

(13 marks)

- (b) Differentiate  $y = \tan^{-1} \left( \frac{1-x}{1+x} \right)$  with respect to  $x$ .

(7 marks)

## Question 6

- (a) Evaluate  $\int_1^2 (x+1)\sqrt{2-x} dx$  by using a suitable substitution or otherwise.

(7 marks)

- (b) (i) Express  $f(x) = \frac{x^2 + 7x + 2}{(1+x^2)(2-x)}$  in terms of partial fractions.

(6 marks)

- (ii) Hence, prove that  $\int_0^1 f(x) dx = \frac{11}{2} \ln 2 - \frac{\pi}{4}$ .

(7 marks)

END OF QUESTION

## APPENDIX 1

## Table of Differentiation

Trigonometric Functions - GENERAL FORM
$\frac{d}{dx}(\sin f(x)) = f'(x) \cos f(x)$
$\frac{d}{dx}(\cos f(x)) = -f'(x) \sin f(x)$
$\frac{d}{dx}(\tan f(x)) = f'(x) \sec^2 f(x)$
$\frac{d}{dx}(\csc f(x)) = -f'(x) \csc f(x) \cot f(x)$
$\frac{d}{dx}(\sec f(x)) = f'(x) \sec f(x) \tan f(x)$
$\frac{d}{dx}(\cot f(x)) = -f'(x) \csc^2 f(x)$

  

Exponential Function - GENERAL FORM
$\frac{d}{dx}(e^{f(x)}) = f'(x) e^{f(x)}$

  

Logarithmic Function - GENERAL FORM
$\frac{d}{dx}(\ln f(x)) = \frac{1}{f(x)} f'(x)$

## APPENDIX 2

Derivatives of Inverse Trigonometric Functions
$\frac{d}{dx}(\sin^{-1} u) = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}, \quad  u  < 1$
$\frac{d}{dx}(\cos^{-1} u) = \frac{-1}{\sqrt{1-u^2}} \frac{du}{dx}, \quad  u  < 1$
$\frac{d}{dx}(\tan^{-1} u) = \frac{1}{1+u^2} \frac{du}{dx}$
$\frac{d}{dx}(\csc^{-1} u) = \frac{-1}{ u \sqrt{u^2-1}} \frac{du}{dx}, \quad  u  > 1$
$\frac{d}{dx}(\sec^{-1} u) = \frac{1}{ u \sqrt{u^2-1}} \frac{du}{dx}, \quad  u  > 1$
$\frac{d}{dx}(\cot^{-1} u) = \frac{-1}{1+u^2} \frac{du}{dx}$

Derivatives of Hyperbolic Functions
$\frac{d}{dx}(\sinh u) = \cosh u \frac{du}{dx}$
$\frac{d}{dx}(\cosh u) = \sinh u \frac{du}{dx}$
$\frac{d}{dx}(\tanh u) = \operatorname{sech}^2 u \frac{du}{dx}$
$\frac{d}{dx}(\operatorname{csch} u) = -\operatorname{csch} u \operatorname{coth} u \frac{du}{dx}$
$\frac{d}{dx}(\operatorname{sech} u) = -\operatorname{sech} u \tanh u \frac{du}{dx}$
$\frac{d}{dx}(\operatorname{coth} u) = -\operatorname{csch}^2 u \frac{du}{dx}$

## APPENDIX 3

Derivatives of Inverse Hyperbolic Functions	
$\frac{d}{dx}(\sinh^{-1} u) = \frac{1}{\sqrt{1+u^2}} \frac{du}{dx}$	
$\frac{d}{dx}(\cosh^{-1} u) = \frac{1}{\sqrt{u^2-1}} \frac{du}{dx}, \quad u > 1$	
$\frac{d}{dx}(\tanh^{-1} u) = \frac{1}{1-u^2} \frac{du}{dx}, \quad  u  < 1$	
$\frac{d}{dx}(\operatorname{csch}^{-1} u) = \frac{-1}{ u \sqrt{1+u^2}} \frac{du}{dx}, \quad u \neq 0$	
$\frac{d}{dx}(\operatorname{sech}^{-1} u) = \frac{-1}{u\sqrt{1-u^2}} \frac{du}{dx}, \quad 0 < u < 1$	
$\frac{d}{dx}(\operatorname{coth}^{-1} u) = \frac{1}{1-u^2} \frac{du}{dx}, \quad  u  > 1$	



## APPENDIX 4

## Trigonometric Identities and Formulas

## FUNDAMENTAL IDENTITIES

$$\csc\theta = \frac{1}{\sin\theta}$$

$$\sec\theta = \frac{1}{\cos\theta}$$

$$\cot\theta = \frac{1}{\tan\theta} = \frac{\cos\theta}{\sin\theta}$$

$$\tan\theta = \frac{\sin\theta}{\cos\theta}$$

$$\sin^2\theta + \cos^2\theta = 1$$

$$1 + \tan^2\theta = \sec^2\theta$$

$$1 + \cot^2\theta = \csc^2\theta$$

## FORMULAS FOR NEGATIVES

$$\sin(-\theta) = -\sin\theta$$

$$\cos(-\theta) = \cos\theta$$

$$\tan(-\theta) = -\tan\theta$$

$$\csc(-\theta) = -\csc\theta$$

$$\sec(-\theta) = \sec\theta$$

$$\cot(-\theta) = -\cot\theta$$

## ADDITION FORMULAS

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

## SUBTRACTION FORMULAS

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

## HALF-ANGLE FORMULAS

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos\theta}{2}}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos\theta}{2}}$$

$$\tan \frac{\theta}{2} = \frac{1 - \cos\theta}{\sin\theta} = \frac{\sin\theta}{1 + \cos\theta}$$

## DOUBLE-ANGLE FORMULAS

$$\sin 2\theta = 2\sin\theta \cos\theta$$

$$\cos 2\theta = \cos^2\theta - \sin^2\theta$$

$$\dots\dots\dots = 1 - 2\sin^2\theta$$

$$\dots\dots\dots = 2\cos^2\theta - 1$$

$$\tan 2\theta = \frac{2\tan\theta}{1 - \tan^2\theta}$$

## PRODUCT-TO-SUM FORMULAS

$$\sin\alpha \sin\beta = \frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

$$\cos\alpha \cos\beta = \frac{1}{2} [\cos(\alpha + \beta) + \cos(\alpha - \beta)]$$

$$\sin\alpha \cos\beta = \frac{1}{2} [\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

$$\cos\alpha \sin\beta = \frac{1}{2} [\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

## SUM-TO-PRODUCT FORMULAS

$$\sin\alpha + \sin\beta = 2\sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$\sin\alpha - \sin\beta = 2\cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

$$\cos\alpha + \cos\beta = 2\cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$\cos\alpha - \cos\beta = -2\sin \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

## APPENDIX 5

## Table of Integration

Trigonometric Functions - GENERALFORM
Where : $f(x) = ax + b$
$\int \cos f(x) dx = \frac{\sin f(x)}{f'(x)} + C$
$\int \sin f(x) dx = \frac{-\cos f(x)}{f'(x)} + C$
$\int \sec^2 f(x) dx = \frac{\tan f(x)}{f'(x)} + C$
$\int \sec f(x) \tan f(x) dx = \frac{\sec f(x)}{f'(x)} + C$
$\int \csc f(x) \cot f(x) dx = \frac{-\csc f(x)}{f'(x)} + C$
$\int \csc^2 f(x) dx = \frac{-\cot f(x)}{f'(x)} + C$
Exponential Function - GENERALFORM
Where : $f(x) = ax + b$
$\int e^{f(x)} dx = \frac{e^{f(x)}}{f'(x)} + C$
Logarithmic Function - GENERALFORM
Where : $f(x) = ax + b$
$\int \frac{1}{f(x)} dx = \frac{\ln  f(x) }{f'(x)} + C$

## APPENDIX 6

Integration of Inverse Trigonometric Functions
$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + C, \quad  x  < a$
$\int \frac{-1}{\sqrt{a^2 - x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + C, \quad  x  < a$
$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C$
$\int \frac{-1}{ x \sqrt{x^2 - a^2}} dx = \frac{1}{a} \csc^{-1}\left(\frac{x}{a}\right) + C, \quad  x  > a$
$\int \frac{1}{ x \sqrt{x^2 - a^2}} dx = \frac{1}{a} \sec^{-1}\left(\frac{x}{a}\right) + C, \quad  x  > a$
$\int \frac{-1}{a^2 + x^2} dx = \frac{1}{a} \cot^{-1}\left(\frac{x}{a}\right) + C$

Integration of Hyperbolic Functions
Where $f(x) = ax + b$ and $f'(x) = a$
$\int \cosh f(x) dx = \frac{\sinh f(x)}{f'(x)} + C$
$\int \sinh f(x) dx = \frac{\cosh f(x)}{f'(x)} + C$
$\int \operatorname{sech}^2 f(x) dx = \frac{\tanh f(x)}{f'(x)} + C$
$\int \operatorname{csch}^2 f(x) dx = \frac{-\operatorname{coth} f(x)}{f'(x)} + C$
$\int \operatorname{sech} f(x) \tanh f(x) dx = \frac{-\operatorname{sech} f(x)}{f'(x)} + C$
$\int \operatorname{csch} f(x) \operatorname{coth} f(x) dx = \frac{-\operatorname{csch} f(x)}{f'(x)} + C$