



**UNIVERSITI KUALA LUMPUR
Malaysia France Institute**

**FINAL EXAMINATION
JULY 2010 SESSION**

SUBJECT CODE : FKB 24302
SUBJECT TITLE : ENGINEERING MATHEMATICS 3
LEVEL : BACHELOR
TIME / DURATION : 4.00 pm – 6.00 pm
(2 HOURS)
DATE : 18 NOVEMBER 2010

INSTRUCTIONS TO CANDIDATES

1. Please read the instructions given in the question paper CAREFULLY.
2. This question paper is printed on both sides of the paper.
3. Please write your answers on the answer booklet provided.
4. Answer should be written in blue or black ink except for sketching, graphic and illustration.
5. This question paper consists of FIVE (5) questions. You are required to answer FOUR (4) questions only. Answer all questions in English.
6. Calculus Formula and Polar graph paper are appended.

THERE ARE 5 PAGES OF QUESTIONS AND 6 PAGES OF APPENDICES, EXCLUDING THIS PAGE.

(Total : 60 marks)

INSTRUCTION : Answer FOUR (4) questions only.

Please use the answer booklet provided.

Question 1

- (a) In the transformation of two dimensional space, find the matrix operator for a reflection on the line $3y = -4x$. (5 marks)

- (b) The transformation L maps the point $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$ to the point $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$ where

$$a = x + 4y + 5z$$

$$b = -4x + 2z$$

$$c = 3z$$

State L , the matrix operator of this transformation.

(2 marks)

- (c) The transformation K has matrix operator $K = \begin{pmatrix} 2 & 0 & -1 \\ 0 & 2 & 1 \\ -3 & -1 & 1 \end{pmatrix}$.

Given that M is the compound transformation consisting of K followed by L from part (b).

- (i) Find the matrix operator, M for this compound transformation. (5 marks)

- (ii) Determine the image of point $A = \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}$ under the transformation M .

(3 marks)

Question 2

- (a) Given $T = 5p^2 + \frac{3}{p}$, determine $\frac{dT}{dq}$ when $p = 2$ given that for that value of p , $\frac{dp}{dq} = 4$

(4 marks)

- (b) Determine the rectangular equation of the following parametric form:

$$x = \frac{1}{t-1}, \quad y = \frac{1}{t^2+1}$$

(4 marks)

- (c) The line segment of the parametric equation $x = \sin^2 t$ and $y = \cos^2 t$ over the interval $0 \leq t \leq \frac{\pi}{2}$ is revolved about the x-axis to generate a cone. Calculate the surface area of the cone.

(7 marks)

Formulas:

$$A = \int_a^b \frac{1}{2} r^2 d\theta$$

$$A = 2\pi \int_A^B x \left\{ \left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right\}^{1/2} dt$$

$$A = 2\pi \int_A^B y \left\{ \left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 \right\}^{1/2} dt$$

Question 3

- (a) Sketch, on the same diagram from **APPENDIX 5**, the polar curves

Circle: $r = 3 \cos \theta$

Cardioid: $r = 1 + \cos \theta$

(5 marks)

- (b) Determine the points of intersection of the circle and the cardioid.

(3 marks)

- (c) Shade the region which lies inside the cardioid and outside the circle and hence calculate the area of the shaded region.

(7 marks)

Question 4

(a) By referring to the Figure 1 below, given that $\vec{OA} = 10\vec{a}$, $\vec{OB} = 8\vec{b}$,

$$\vec{DB} = \frac{2}{3}\vec{OB} \text{ and } \vec{BC} = \frac{2}{3}\vec{CA}$$




Figure 1

Determine in terms of \vec{a} and \vec{b} , the following vectors

(i) \vec{BA} (2 marks)

(ii) \vec{BC} (4marks)

(iii) \vec{DC} (4marks)

(b) A firefly is flying in still air at 205.5 km/h on a bearing of 45.7° . A steady wind suddenly springs up, blowing due south at 67.3 km/h. By using the Cosine Law, find the velocity of the firefly over the ground.

(5 marks)

Question 5

(a) Given $\vec{OA} = 3i + 4j$ and $\vec{OB} = 2hi - hj$.

(i) Determine \vec{AB} in terms of h .

(1 mark)

(ii) If $\vec{AB} = 9i + kj$, calculate the value of h and k .

(2 marks)

(b) Given that $\vec{OP} = 2i + j - 3k$, $\vec{OQ} = i - 2j + k$ and $\vec{OR} = -i + j - 4k$.

Determine the following:

(i) $\vec{OP} \cdot (\vec{OQ} \times \vec{OR})$

(2 marks)

(ii) $\vec{OQ} \times \frac{1}{2} \vec{OP}$

(2 marks)

(iii) $\left| \vec{OQ} \times \frac{1}{2} \vec{OP} \right|$

(2 marks)

(c) A triangle ABC has its vertices at the points $A(3, -1, 4)$, $B(1, 5, -4)$ and $C(-6, 2, 2)$.

Obtain, in the form of $ai + bj + ck$, the vectors represent \vec{AB} and \vec{CA} .

(6 marks)

END OF QUESTION

APPENDIX 1

STANDARD TRANSFORMATIONS

(1)	Rotation through an angle , θ , about the origin	$M = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$
(2)	Rotation through $\frac{\pi}{2}$ clockwise about the origin	$M = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$
(3)	Rotation through $\frac{\pi}{2}$ anti-clockwise about the origin	$M = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$
(4)	Reflection in the x-axis	$M = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$
(5)	Reflection in the y-axis	$M = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$
(6)	Reflection in the line : $y = x$ or $y - x = 0$	$M = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
(7)	Reflection in the line : $y = -x$ or $y + x = 0$	$M = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$
(8)	Reflection in the line : $y = mx$ or $y = x \tan \theta$ NOTE : $m = \tan \theta$	$M = \begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix}$ where :

		$\cos 2\theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - m^2}{1 + m^2}$ $\sin 2\theta = \frac{2 \tan \theta}{1 + \tan^2 \theta} = \frac{2m}{1 + m^2}$
(9)	Shear of θ in the direction O_x	$M = \begin{pmatrix} 1 & \tan \theta \\ 0 & 1 \end{pmatrix}$
(10)	Shear of θ in the direction O_y	$M = \begin{pmatrix} 1 & 0 \\ \tan \theta & 1 \end{pmatrix}$

APPENDIX 2

TRIGONOMETRIC IDENTITIES AND FORMULAS

FUNDAMENTAL IDENTITIES	FORMULAS FOR NEGATIVES
$\csc \theta = \frac{1}{\sin \theta}$	$\sin(-\theta) = -\sin \theta$
$\sec \theta = \frac{1}{\cos \theta}$	$\cos(-\theta) = \cos \theta$
$\cot \theta = \frac{1}{\tan \theta} = \frac{\cos \theta}{\sin \theta}$	$\tan(-\theta) = -\tan \theta$
$\tan \theta = \frac{\sin \theta}{\cos \theta}$	$\csc(-\theta) = -\csc \theta$
$\sin^2 \theta + \cos^2 \theta = 1$	$\sec(-\theta) = \sec \theta$
$1 + \tan^2 \theta = \sec^2 \theta$	$\cot(-\theta) = -\cot \theta$
$1 + \cot^2 \theta = \csc^2 \theta$	
ADDITION FORMULAS	SUBTRACTION FORMULAS
$\sin(A + B) = \sin A \cos B + \cos A \sin B$	$\sin(A - B) = \sin A \cos B - \cos A \sin B$
$\cos(A + B) = \cos A \cos B - \sin A \sin B$	$\cos(A - B) = \cos A \cos B + \sin A \sin B$
$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$	$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$
HALF-ANGLE FORMULAS	DOUBLE-ANGLE FORMULAS
$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$	$\sin 2\theta = 2 \sin \theta \cos \theta$
$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$	$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$ = $1 - 2 \sin^2 \theta$ = $2 \cos^2 \theta - 1$
$\tan \frac{\theta}{2} = \frac{1 - \cos \theta}{\sin \theta} = \frac{\sin \theta}{1 + \cos \theta}$	$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$
PRODUCT-TO-SUM FORMULAS	SUM-TO-PRODUCT FORMULAS
$\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)]$	$\sin \alpha + \sin \beta = 2 \sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$
$\cos \alpha \sin \beta = \frac{1}{2} [\sin(\alpha + \beta) - \sin(\alpha - \beta)]$	$\sin \alpha - \sin \beta = 2 \cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$
$\cos \alpha \cos \beta = \frac{1}{2} [\cos(\alpha + \beta) + \cos(\alpha - \beta)]$	$\cos \alpha + \cos \beta = 2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$
$\sin \alpha \sin \beta = \frac{1}{2} [\cos(\alpha - \beta) - \cos(\alpha + \beta)]$	$\cos \alpha - \cos \beta = -2 \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$

APPENDIX 3

DIFFERENTIATION

TRIGONOMETRIC FUNCTIONS

STANDARD FORM	GENERAL FORM
$\frac{d}{dx}(\sin x) = \cos x$	$\frac{d}{dx}(\sin f(x)) = \cos f(x) f'(x)$
$\frac{d}{dx}(\cos x) = -\sin x$	$\frac{d}{dx}(\cos f(x)) = -\sin f(x) f'(x)$
$\frac{d}{dx}(\tan x) = \sec^2 x$	$\frac{d}{dx}(\tan f(x)) = \sec^2 f(x) \cdot f'(x)$
$\frac{d}{dx}(\csc x) = -\csc x \cot x$	$\frac{d}{dx}(\csc f(x)) = -\csc f(x) \cot f(x) \cdot f'(x)$
$\frac{d}{dx}(\sec x) = \sec x \tan x$	$\frac{d}{dx}(\sec f(x)) = \sec f(x) \tan f(x) \cdot f'(x)$
$\frac{d}{dx}(\cot x) = -\csc^2 x$	$\frac{d}{dx}(\cot f(x)) = -\csc^2 f(x) \cdot f'(x)$

EXPONENTIAL FUNCTION

STANDARD FORM	GENERAL FORM
$\frac{d}{dx}e^x = e^x$	$\frac{d}{dx}e^{f(x)} = e^{f(x)} \cdot f'(x)$

LOGARITHMIC FUNCTION

STANDARD FORM	GENERAL FORM
$\frac{d}{dx} \ln x = \frac{1}{x}$	$\frac{d}{dx} \ln f(x) = \frac{f'(x)}{f(x)}$

APPENDIX 4

INTEGRATION

TRIGONOMETRIC FUNCTIONS

STANDARD FORM	GENERAL FORM Where : $f(x) = ax + b$
$\int \cos x dx = \sin x + c$	$\int \cos f(x) dx = \frac{\sin f(x)}{f'(x)} + c$
$\int \sin x dx = -\cos x + c$	$\int \sin f(x) dx = \frac{-\cos f(x)}{f'(x)} + c$
$\int \sec^2 x dx = \tan x + c$	$\int \sec^2 f(x) dx = \frac{\tan f(x)}{f'(x)} + c$
$\int \sec x \tan x dx = \sec x + c$	$\int \sec f(x) \tan f(x) dx = \frac{\sec f(x)}{f'(x)} + c$
$\int \csc x \cot x dx = -\csc x + c$	$\int \csc f(x) \cot f(x) dx = \frac{-\csc f(x)}{f'(x)} + c$
$\int \csc^2 x dx = -\cot x + c$	$\int \csc^2 f(x) dx = \frac{-\cot f(x)}{f'(x)} + c$

EXPONENTIAL FUNCTION

STANDARD FORM	GENERAL FORM Where : $f(x) = ax + b$
$\int e^x dx = e^x + c$	$\int e^{f(x)} dx = \frac{e^{f(x)}}{f'(x)} + c$

FORM OF 1/x

STANDARD FORM	GENERAL FORM Where : $f(x) = ax + b$
$\int \frac{1}{x} dx = \ln x + c$	$\int \frac{1}{f(x)} dx = \frac{\ln f(x) }{f'(x)} + c$

APPENDIX 5

NAME :

ID NUMBER :

